

Patient Name : Demo Patient Name
Age / Sex : 41 Y / M
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 19-Jan-25 11:34
Patient ID : UHID.DEMO.001

H Pylori Antigen (Rapid Stool)

Stool Sample

Accession No: DEMO_BARCODE **Collected On:** 19-Jan-25 11:34 **Received On:** 19-Jan-25 14:04 **Approved On:** 19-Jan-25 17:19

Observation	Result	Unit	Biological Ref. Interval	Method
Helicobacter Pylori Antigen	Positive			Immunochromatography

Interpretation

- Positive - Indicates presence of Helicobacter pylori Antigen
- Negative - Indicates absence of Helicobacter pylori Antigen

Note

- This is a qualitative test.
- Positive results in stool are seen in gastrointestinal diseases of stomach and duodenum like Gastritis, Dyspepsia and peptic ulcer disease caused by Helicobacter pylori (H pylori).
- Negative results are seen in absence of H pylori infection. However, it does not rule out the disease.
- Results should be correlated with clinical conditions and other tests like Culture and Endoscopy.
- False-positive reaction could be a consequence of cross-reaction with other helicobacter species or other microorganisms of intestinal flora.
- False negative reaction may be due to processing of sample collected after ingestion of antimicrobials, proton pump inhibitors and bismuth preparations for treatment as they are known to suppress H pylori. The test should be repeated on a new specimen obtained two weeks after discontinuing treatment.

Uses

- To establish the presence and possible etiologic role of H pylori in cases of chronic gastric ulcer, gastritis, duodenal ulcer, dyspepsia, etc.
- To evaluate efficacy of treatment.

Advise: Please correlate result clinically.



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

