

Patient Name : Demo Patient Name
Age / Sex : 48 Y / F
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 21-Jan-25 18:36
Patient ID : UHID.DEMO.001

RA Factor				Serum Sample
Accession No: DEMO_BARCODE		Collected On: 21-Jan-25 18:36	Received On: 21-Jan-25 19:24	Approved On: 21-Jan-25 20:56
Observation	Result	Unit	Biological Ref. Interval	Method
RA Factor [Quantitative]	<8.6	IU/ml	<12.0	Turbidimetry

Technology: VITROS MicroTip, MicroSensor and Intellicheck Technology
Analyzer: Fully Automated Integrated Biochemistry and Immunology Analyzer: VITROS 5600

Clinical Significance: Rheumatoid factor (RF) consists of autoantibodies of immunoglobulin isotype IgM, IgA, IgG and IgE. The function of RF remains unclear, but it may play a role in the regulation of humoral and cellular immunity and protection against invading microorganisms. Most patients with rheumatoid arthritis and Sjogren's syndrome have elevated levels of RF. RF may also be elevated in scleroderma, dermatomyositis, Waldenström's disease, sarcoidosis, and systemic lupus erythematosus. There are also instances of elevated RF without apparent disease or definable clinical disorders.

Remarks: Please correlate results with clinical conditions.



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

