

Patient Name : Demo Patient Name
Age / Sex : 10 Y / M
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 21-Jan-25 12:36
Patient ID : UHID.DEMO.001

AntiStreptolysin O

Serum Sample

Accession No: DEMO_BARCODE **Collected On:** 21-Jan-25 12:36 **Received On:** 22-Jan-25 08:42 **Approved On:** 22-Jan-25 10:41

Observation	Result	Unit	Biological Ref. Interval	Method
Anti-Streptolysin O	<25	IU/mL	0 - 240 IU/mL	Latex Agglutination

Clinical Notes:

* ASO titre is used to assist in the diagnosis of rheumatic fever, scarlet fever and post-streptococcal glomerulonephritis.
* A titre has a significance only if it is greatly elevated, or if a rise in titre can be demonstrated in paired blood samples taken days apart.* The antibody levels begin to rise after 1 to 3 weeks of strep infection, peaks in 3 to 5 weeks and falls back to insignificant levels in 6 months.
* False positive results may be seen in patients suffering from tuberculosis and liver disease.
* In case of suspected false negative result, anti-DNase titre is advisable.

Please correlate results clinically.



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

