

Patient Name : Demo Patient Name
Age / Sex : 1 Y 1 M 20 D / M
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 21-Jan-25 10:19
Patient ID : UHID.DEMO.001

Anti HAV IgM

Serum Sample

Accession No: DEMO_BARCODE **Collected On:** 21-Jan-25 10:19 **Received On:** 21-Jan-25 13:43 **Approved On:** 21-Jan-25 18:02

Observation	Result	Unit	Biological Ref. Interval	Method
Anti Hepatitis A Virus (Anti-HAV) - IgM	0.30		<0.90	CLIA

Clinical Significance of Anti HAV IgM

The route of HAV infection is predominantly oral-fecal with an incubation period of 2-7 weeks, during which HAV can be detected in stools. HAV infection does not originate any chronic hepatitis and complications are uncommon. The infection stimulates a strong immunological response in the patient with elevated titres first of IgM and then IgG, whose final presence lasts for years after infection.

Remarks: Please correlate results with clinical conditions.



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

