

Patient Name : Demo Patient Name
Age / Sex : 30 Y / F
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 19-Jan-25 08:31
Patient ID : UHID.DEMO.001

Anti Cardiolipin Antibody [IgM]

Serum Sample

Accession No: DEMO_BARCODE **Collected On:** 19-Jan-25 08:31 **Received On:** 19-Jan-25 11:19 **Approved On:** 19-Jan-25 14:45

Observation	Result	Unit	Biological Ref. Interval	Method
Anti Cardiolipin Antibody [IgM]	3.6	U/mL	<20	CLIA

Biological Reference Range for Anti Cardiolipin [IgM]:

- Negative: Less Than 20 U/mL
- Intermediate (Borderline): 20-30 U/mL
- Positive : More Than 30 U/mL

Technology: Test Performed on Fully Automated - ACL Acstar - CLIA Analyzer

2006 International Consensus statement on Classification of definite APS

Clinical Criteria	Lab Criteria
Arterial/Venous thrombosis	Cardiolipin antibodies (aCL)
Fetal Loss	Beta 2 Glycoprotein 1 antibodies
Premature birth	Lupus Anticoagulant (LA)

Clinical Significance of Cardiolipin [IgM] Levels:

Measurement of cardiolipin antibodies is the standard procedure for the detection of suspected antiphospholipid syndrome (APS). High aCL concentrations are associated with increased risk of venous and arterial thrombosis, recurrent pregnancy loss and thrombocytopenia.

Remarks: Please correlate results with clinical conditions.



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.



Scan to Validate