

Patient Name : Demo Patient Name
Age / Sex : 26 Y / M
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 11-Aug-25 11:23
Patient ID : UHID.DEMO.001

HSV-II IgG (Herpes Simplex Virus)

Serum Sample

Accession No: DEMO_BARCODE Collected On: 11-Aug-25 11:23 Received On: 11-Aug-25 11:49 Approved On: 11-Aug-25 15:30

Observation	Result	Unit	Biological Ref. Interval	Method
Herpes Simplex Virus (HSV) - II IgG	<1	AU/mL	<9:Non-Reactive <9-13:Equivocal ≥13:Reactive	CLIA

Clinical Significance

- A **non-reactive** result indicates that the specific immunity has not been acquired but cannot rule out the early stage of acute infection.
- A **reactive** result indicates either early acute infection or past exposure to the pathogen. If it is suspected to be early acute infection, a HSV-1 IgM test or other serological method for detection of additional HSV-2 markers, such as a HSV-2 IgG Avidity test, could be performed.
- For **equivocal** result, a second sample should be taken and repeated HSV-2 IgG testing no less than one or two weeks later, or/and using a HSV-2 IgM test to confirm.
- Diagnosis of infectious diseases **should not be established on the basis of a single test result**, but should be determined in conjunction with clinical findings and other diagnostic procedures as well as in association with medical judgment.

Remarks: Please correlate with clinical conditions



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

