

Patient Name : Demo Patient Name
Age / Sex : 26 Y / M
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 11-Aug-25 11:23
Patient ID : UHID.DEMO.001

HSV-II IgM (Herpes Simplex Virus)

Serum Sample

Accession No: DEMO_BARCODE **Collected On:** 11-Aug-25 11:23 **Received On:** 11-Aug-25 11:49 **Approved On:** 11-Aug-25 15:30

Observation	Result	Unit	Biological Ref. Interval	Method
Herpes Simplex Virus (HSV) - II IgM	<1	AU/mL	<6:Non-Reactive <6-10:Equivocal ≥10:Reactive	CLIA

Clinical Significance

- A **non-reactive** result cannot always rule out acute HSV-2 infection, because the infection maybe be in its very early stage and the patient is still unable to synthesize HSV-2 specific IgM.
- An **equivocal** result may be indicative either of recent infection or of past infection with long-lasting HSV-2 IgM. A second sample should be collected within a reasonable period of time (e.g., within one week). Serological data from detection of additional HSV-1 markers may provide useful information for clinical interpretation of results.
- Diagnosis of infectious diseases should not be established on the basis of a single test result, but should be determined in conjunction with clinical findings and other diagnostic procedures as well as in association with medical judgment.

Remarks: Please correlate with clinical conditions



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

