

Patient Name : Demo Patient Name
Age / Sex : 30 Y / F
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 19-Jan-25 08:31
Patient ID : UHID.DEMO.001

Beta-2 Glycoprotein IgG

Serum Sample

Accession No: DEMO_BARCODE Collected On: 19-Jan-25 08:31 Received On: 19-Jan-25 11:19 Approved On: 19-Jan-25 14:45

Observation	Result	Unit	Biological Ref. Interval	Method
Beta-2 Glycoprotein I IgG	< 6.4	U/mL	<20	CLIA

Reference Range:

Negative: <20
Positive: >20

Technology: Test Performed on Fully Automated - ACL Acustar - CLIA Analyzer

Clinical Significance: This test screens for the presence of IgG antibodies to Beta-2 Glycoprotein which are associated with arterial and venous thrombosis and recurrent abortions.

2006 International Consensus statement on Classification of definite APS

Clinical Criteria	Lab Criteria
Arterial/Venous thrombosis	Cardiolipin antibodies (aCL)
Fetal Loss	Beta 2 Glycoprotein 1 antibodies
Premature birth	Lupus Anticoagulant (LA)

Indications:

- Clinical history suggestive of Arterial or venous thrombosis.
- History of Recurrent fetal loss/Intrauterine growth restriction/ premature birth.
- Thrombocytopenia.

Advise: Please correlate results clinically.



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

