

Patient Name : Demo Patient Name
Age / Sex : 24 Y / F
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 20-Jan-25 11:54
Patient ID : UHID.DEMO.001

Toxoplasma [IgM]				Serum Sample
Accession No: DEMO_BARCODE		Collected On: 20-Jan-25 11:54	Received On: 20-Jan-25 15:13	Approved On: 20-Jan-25 16:52
Observation	Result	Unit	Biological Ref. Interval	Method
Toxoplasma [IgM]	<1	AU/mL	<6:Non-Reactive <6-10:Equivocal ≥10:Reactive	CLIA

Clinical Notes:

- A non-reactive result indicates that the specific immunity has not been acquired but cannot rule out the early stage of acute infection. Patients with non-reactive results with suspected exposure to the Toxoplasma gondii virus should be retested within 2 weeks.
- A reactive result indicates either early acute infection or past exposure to the pathogen. If it is suspected to be early acute infection, a second test or other serological method for detection of additional Toxoplasma gondii markers could be performed.
- For equivocal result, a second sample should be taken and repeated Toxo IgM testing no less than one or two weeks later.
- Diagnosis of infectious diseases should not be established on the basis of a single test result, but should be determined in conjunction with clinical findings and other diagnostic procedures as well as in association with medical judgment.

Remarks: Please correlate results with clinical conditions and drug history.



This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

