

Patient Name : Demo Patient Name
Age / Sex : 29 Y / F
Referred By : DEMO HOSPITAL
Centre : HOD Head Office

Lab No : Demo Visit No
Registration On : 21-Jan-25 17:00
Patient ID : UHID.DEMO.001

Pregnancy Test [Urine]				Urine Sample
Accession No: DEMO_BARCODE		Collected On: 21-Jan-25 17:00	Received On: 21-Jan-25 17:18	Approved On: 21-Jan-25 17:23
Observation	Result	Unit	Biological Ref. Interval	Method
Pregnancy Test	Positive			Immunochromatography

Clinical Notes: Most doctors recommend that you wait until the first day of your missed period before taking a urine pregnancy test. This is usually about two weeks after conception. Urine pregnancy tests are around 97% accurate when done correctly.

- **Limitations:** A number of conditions other than pregnancy including trophoblastic and non-trophoblastic neoplasms such as Hydatiform mole, choriocarcinoma etc. cause elevated levels of hCG. Such clinical conditions must be ruled out before diagnosis of pregnancy can be made.

- Highly dilute urine specimens and specimens from very early pregnancy may not contain representative levels of hCG. If pregnancy is still suspected, repeat the test with first morning urine 48-72 hours after initial test.

Advised: Correlation with Beta HCG blood tests for better sensitivity. A Beta HCG blood test can generally detect a pregnancy earlier than a urine test, at about 7-12 days from possible conception (but if a negative result is received, a test should be repeated if a period is missed.)

Remark: Please correlate results clinically.

 This is a Demo Signature
and the doctor's signature should appear here

In case of any unexpected or alarming results, please contact us immediately for re-confirmation, clarifications, and rectifications, if needed.

